

Quorum™ Clear V

Disinfectant - Sanitizer - Deodorizer - One-Step Cleaner/Disinfectant for Industrial and Commercial Use

Dairies - Beverage and Food Processing Plants -
Farm Premises - Poultry Premises - Poultry Hatcheries -
Animal Housing Facilities

ACTIVE INGREDIENTS:

Alkyl (C ₁₄ , 50%; C ₁₂ , 40%; C ₁₆ , 10%)	
Dimethyl benzyl ammonium chloride.....	3.00%
Octyl decyl dimethyl ammonium chloride	2.25%
Didecyl dimethyl ammonium chloride	1.35%
Dioctyl dimethyl ammonium chloride	0.90%
OTHER INGREDIENTS:	92.50%
TOTAL:	100.00%

KEEP OUT OF REACH OF CHILDREN

DANGER

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER: CORROSIVE: Causes irreversible eye damage and skin burns. Do not get in eyes, on skin or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses) protective clothing and rubber gloves. Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

DO NOT MIX WITH ANYTHING BUT WATER

EPA Reg. No. 1677-198

Read other label for directions for use.

764261/5301/0116

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

FOR EMERGENCY MEDICAL INFORMATION IN USA OR CANADA, CALL: 1-800-328-0026.

FOR EMERGENCY MEDICAL INFORMATION WORLDWIDE, CALL: 1-651-222-5352 (IN THE USA).

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.


STORAGE & DISPOSAL

DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL.

PESTICIDE STORAGE: Store in original container in areas inaccessible to children. Do not store on side. Avoid creasing or impacting of side walls.


PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING AND DISPOSAL: Non-refillable container. Do not reuse or refill this container. Triple rinse as follows: Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Follow pesticide disposal instructions for rinsate disposal. Repeat procedure two more times. Then offer for recycling or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by incineration.



Disinfectant/Sanitizer

NET CONTENTS:
55 US GAL (208 L)
6300567





Quorum™ Clear V

Disinfectant - Sanitizer - Deodorizer - One-Step Cleaner/Disinfectant for Industrial and Commercial Use

KEEP OUT OF REACH OF CHILDREN DANGER

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish and aquatic invertebrates. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Apply Quorum Clear V with a cloth, mop or mechanical spray device. When applied with a mechanical spray device, surface must be sprayed until thoroughly wetted. For sprayer applications, use a coarse spray device. Spray 6-8 inches from surface; rub with brush, sponge or cloth. Do not breathe spray. Fresh solution must be prepared daily or when the use solution becomes visibly dirty.

GENERAL DISINFECTION/VIRUCIDAL* DIRECTIONS: At 1 oz. per 1 gallon of water, this product is an effective general disinfectant that can be used to disinfect floors, walls, finished woodwork, sink tops, garbage pails, telephones, rest rooms, bathroom fixtures, and other hard, nonporous surfaces. For heavily soiled areas, a pre-cleaning step is required. Apply disinfecting solution to surface by flushing, mopping, sponging or hand pump spraying so as to thoroughly wet surface. Allow surface to remain wet for 10 minutes. Remove solution with a clean mop, cloth, or wet vacuum pickup or allow to air dry. Prepare fresh solution at least daily or when use solution becomes visibly dirty. Any food contact surfaces that have been disinfected must be thoroughly rinsed with potable water. Do not use this product as a disinfectant on glasses, dishes or utensils.

Before use in federally inspected meat and poultry food processing plants and dairies, food products and packaging materials must be removed from the room or carefully protected.

Quorum Clear V is effective against: *Avian Influenza Virus and *Newcastle's Disease Virus.

Poultry and Swine Premise Disinfection:

1. Remove all poultry and feeds from premises, vehicles and enclosures such as coops and crates.
2. Remove all litter, droppings and manure from floors, walls and surfaces of barns, pens, stalls, chutes and other facilities and fixtures occupied or traversed by animals.
3. Empty all troughs, racks and other feeding and watering appliances.
4. Thoroughly clean all surfaces with a detergent and rinse with water.
5. Saturate surfaces with 4.0 oz. per 4 gallons of water Quorum Clear V solution for a period of at least 10 minutes. Wipe or allow to air dry.
6. Immerse all halters, ropes and other types of equipment used in handling and restraining animals, as well as forks, shovels and scrapers used for removing litter and manure. Wipe or allow to air dry.
7. Ventilate buildings, cars, trucks, coops and other closed spaces. Do not house poultry or employ equipment until treatment has been absorbed, set or dried.
8. Thoroughly scrub treated feed racks, troughs, automatic feeders, fountains and waterers with a detergent and rinse with potable water before reuse.

See your Ecolab representative for specific recommendations for all cleaning and rinsing requirements.

Poultry Hatchery Disinfection: Use to treat hatchers, setters, trays, racks, carts, sexing tables, delivery trucks and other hard non-porous surfaces. Clean out any remaining eggs and chicks. Remove gross soils, such as litter, down, shell fragments or other hatching related debris. Empty all racks and other equipment. Thoroughly

wash all surfaces, including floors, walls, conveyors, trays and water systems with a recommended detergent. Rinse thoroughly with water. Apply 4.0 oz. per 4 gallons of water solution of Quorum Clear V with a mop, cloth, brush or coarse spray. Apply to hatchers, setters, trays, racks, sexing tables, delivery trucks and other hard surfaces. Wet surfaces and allow to remain wet for 10 minutes. Ventilate buildings and other closed spaces. Allow to dry before reintroducing eggs.

Farm Premise Disinfection Directions:

1. Remove all animals and feed from premises, vehicles, and enclosures.
2. Remove all litter, droppings and manure from floors, walls and surfaces of facilities occupied or traversed by animals.
3. Empty all troughs, racks and other feeding and watering appliances.
4. Thoroughly clean surfaces with soap or detergent by manual, foam, or spray application and rinse with water.
5. Saturate surfaces with a use-solution of 4 oz. of Quorum Clear V per 4 gallons of water (or equivalent dilution) and allow to remain wet for a period of at least 10 minutes. Wipe or allow to air dry.
6. Immerse all halters, ropes and other types of equipment used in handling and restraining animals as well as forks, shovels and scrapers used for removing litter and manure. Wipe or allow to air dry.
7. Ventilate buildings and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried.
8. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains and waterers with soap or detergent, and rinse with potable water before reuse.

Deodorizing Directions: This product deodorizes garbage storage areas, garbage bins, toilet bowls and any other hard non-porous surfaces in odor causing areas. Mix 1/2 oz. per gallon of water and apply solution to surfaces. Thoroughly wet surfaces, allow to air dry.

Non-Food Contact Surface Sanitizing: Add 0.341 oz. of Quorum Clear V per 1 gallon of water (200 ppm active quat) to sanitize pre-cleaned, hard, nonporous, non-food contact surfaces against *Staphylococcus aureus* and *Enterobacter aerogenes*. Apply sanitizer use-solution with a cloth, mop, sponge, sprayer or by immersion. For sprayer applications, use a coarse spray device. Spray 6-8 inches from surface. Do not breathe spray. Treated surfaces must remain wet for 5 minutes. Wipe dry with a sponge, mop, or cloth or allow to air dry.

Entryway Sanitizing Systems: To prevent cross contamination from area to area, set the system to deliver sanitizing solution at 0.68 - 2.04 oz. per gallon of water (400-1200 ppm active quat). The foam (or spray) must cover the entire path of the doorway. Set the system so that a continuous wet blanket of sanitizer solution is delivered to the floor. Do not mix other foam additives to the sanitizing solution.

Shoe Bath Sanitizer Directions: To prevent cross contamination into animal areas and the packaging and storage areas of food plants, shoe baths containing one inch of freshly made solution must be placed at all entrances to buildings and hatcheries. Scrape waterproof work boots (shoes) and place in solution of 0.341 oz. of Quorum Clear V per 1 gallon of water (200 ppm) (or equivalent dilution) for 60 seconds prior to entering area. If there is a heavy soil load or excessive traffic, place work boots (shoes) in a use-solution of 0.68 - 1.4 oz. per 1 gallon of water (400 - 800 ppm active quat)(or equivalent dilution) for 60 seconds prior to entering area. Change the solution in the bath daily or more often if solution appears diluted or soiled.

Shoe Foam Sanitizer Directions: Quorum Clear V can be used to prevent cross contamination into animal areas and packaging and storage areas of food plants. Apply a foam layer approximately 0.5 to 2 inches thick made from a solution of 1.364 to 2 ounces per gallon of water (or equivalent use dilution) (800 to 1200 ppm active) at all entrances to buildings, hatcheries, production and packaging rooms by using a foam generating machine or aerator to apply foam layer. Follow the foaming directions as specified by the manufacturer of the foam generator/aerator. Scrape waterproof shoes. Stand and/or walk through foamed area for 60 seconds prior to entering area. Foam area must be washed and replaced daily or when it appears dirty. Quorum Clear V can be used to sanitize hard, non-porous food contact surfaces such as tables, counters, utensils and food processing equipment.

Quorum Clear V is an effective sanitizer against *Escherichia coli* and *Staphylococcus aureus* on food contact surfaces when used at 0.26 - 0.68 oz. per 1 gallon of 400 ppm hard water (150 ppm to 400 ppm active quat). Quorum Clear V is an effective sanitizer against *Escherichia coli*, *Staphylococcus aureus*, *Campylobacter jejuni*, *Escherichia coli* O157:H7, *Klebsiella pneumoniae*, *Listeria monocytogenes*, *Salmonella enterica*, *Shigella sonnei*, *Yersinia enterocolitica* and *Enterobacter sakazakii* on food contact surfaces when used at 0.341 - 0.68 oz. per 1 gallon of 500 ppm hard water (200 ppm to 400 ppm active quat).

To Sanitize Food Contact Surfaces, Food Processing Equipment and Other Hard Surfaces in Food Processing Locations, Dairies, or Restaurants: Use Quorum Clear V to sanitize pre-cleaned hard non-porous surfaces of food processing equipment, dairy equipment, food utensils, dishes, silverware, glasses, sink tops, countertops, refrigerated storage areas and display equipment and other hard non-porous surfaces in federally inspected meat and poultry plants or restaurants. Prior to application, remove gross food particles and soil by a pre-flush, pre-scrape or when necessary, a pre-soak. Then thoroughly wash or flush objects with a good detergent or compatible cleaner followed by a potable water rinse before application of the sanitizing solution. Apply a use-solution of 1.04 oz. to 2.72 oz. of Quorum Clear V per 4 gallons of 400 ppm hard water (150 - 400 ppm active quat)

or 1.364 oz. to 2.72 oz. of Quorum Clear V per 4 gallons of 500 ppm hard water (200 - 400 ppm active quat). Expose all surfaces to the sanitizing solution for a period of not less than 1 minute. Allow equipment to drain thoroughly and air dry.

In dairy processing facilities and restaurants in Wisconsin, clean equipment with a good detergent and follow with a potable water rinse, then rinse equipment with a sanitizing solution of 1.364 oz. to 2.72 oz. of Quorum Clear V per 4 gallons of 500 ppm hard water (200 - 400 ppm active quat).

SANITIZING HARD, NON-POROUS, NON-EDIBLE OUTSIDE SURFACES OF AIRTIGHT, SEALED PACKAGES CONTAINING FOOD OR NON-FOOD PRODUCTS: Quorum Clear V may be used as a final sanitizing rinse for hard, non-porous non-edible outside surfaces of airtight, sealed packages containing food or non-food products at a use-solution of 1.04 oz. to 2.72 oz. of Quorum Clear V per 4 gallons of 400 ppm hard water (150 - 400 ppm active quat) or 1.364 oz. to 2.72 oz. of Quorum Clear V per 4 gallons of 500 ppm hard water (200 - 400 ppm active quat). The treated hard, non-porous, non-edible packaging, such as food wraps and meat casings, must be removed and discarded before packaged food products are further processed or consumed. All surfaces must be exposed to the sanitizing solution for a period of not less than 1 minute. Drain thoroughly. No rinse necessary. This is not to be used on porous surfaces.

For Continuous Treatment of Conveyors: Remove gross food particles and excess soil by a pre-flush or pre-scrape. Wash with a good detergent or compatible cleaner. Rinse equipment thoroughly with potable water, then rinse equipment with a sanitizing solution. During processing, apply Quorum Clear V at a 150 to 400 ppm quat level to conveyors with MIKRO MASTER or other suitable feeding equipment. Controlled volumes of sanitizer are applied to return portion of conveyor through nozzles so located as to permit maximum drainage of sanitizer from equipment and to prevent puddles on top of belt. During interruptions in operation, coarse spray equipment, peelers, collators, slicers and saws with MIKRO MASTER dispensed Quorum Clear V solution of 150 to 400 ppm quat. Conveyors and other equipment must be free of product when applying this coarse spray. Use 200 ppm to 400 ppm quat level in Wisconsin dairy processing facilities.

Sanitizing Shell Eggs Intended For Food: To sanitize previously cleaned food-grade eggs in shell egg and egg product processing plants, spray with a solution of 1.04 oz. to 2.72 oz. of Quorum Clear V in 4 gallons of warm water (150 to 400 ppm). The solution must be warmer than the eggs, but not to exceed 130°F. Wet eggs thoroughly and allow to drain. Eggs sanitized with Quorum Clear V shall be subjected to a potable water rinse only if broken immediately for use in the manufacture of egg products. Eggs must be reasonably dry before casing or breaking. Do not reuse solution for sanitizing eggs.

Note: Only clean, whole eggs can be sanitized. Dirty, cracked or punctured eggs cannot be sanitized.

Sanitizing - Non-Porous Gloved Hands: To prevent cross contamination into animal areas and the packaging and storage areas of food plants, dip, soak or spray pre-washed (plastic, latex or other synthetic rubber) gloved hands so that there is enough sanitizing solution to cover the gloved area. For sprayer applications, use a coarse spray device and spray the glove surfaces until thoroughly wetted. Do not breathe spray. **Do not let sanitizing solution come in contact with exposed skin.** Make up the sanitizing solution by adding 1.04 to 2.72 ounces of product per 4 gallons of water (or equivalent use dilution) (150-400 ppm active quat). Dip, soak or spray in solution and allow gloved hands to remain wet for at least 60 seconds. Allow them to air dry. No potable water rinse is allowed. Change the sanitizing solution at least daily or when solution appears dirty.

Hatchery Rooms: Remove all animals and feed from premises, vehicles and enclosures. Remove all litter and manure from floors, walls, and surfaces of the room to be treated. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Close room off so saturate is confined to the room to be treated. Mix 187 ounces of this product to 2.5 gallons water and apply by dipping, soaking, spraying or flooding (as appropriate) throughout the room to assure the hard surfaces are completely wet. A dust mist respirator must be worn when applying the product. Do not house livestock or employ equipment until treatment has been absorbed or dried. **Do not allow people or animals to contact or breathe this saturate for a minimum of 2 hours.** Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturation is to be used in addition to acceptable manual cleaning and disinfecting of room and machine surfaces.

NOTE:

FOR MECHANICAL OPERATIONS: A prepared use solution may not be reused for sanitizing, but may be reused for other purposes such as cleaning.

FOR MANUAL OPERATIONS: Prepare a fresh sanitizing solution as soon as it becomes diluted or soiled.

EPA Reg. No. 1677-198
EPA Est. 1677-IL-2 (J), 1677-TX-1 (D),
1677-GA-1 (M), 1677-CA-1 (S), 1677-MN-1 (P),
70271-CA-2 (A), 1677-CA-2 (R), 1677-WV-1 (V),
303-IN-1 (L).
Superscript refers to first letter of date code.

Ecolab Inc. · 370 Wabasha Street N ·
St Paul MN 55102-1390 USA ·
tel. 1 800 392 3392
© 2016 Ecolab USA · All rights reserved
Made in United States
734937/5301/0116